



HIV and AIDS in Victoria —1983 to 2004




Newly Acquired HIV = Data collected since 1991. Individuals classified on the basis of a previous negative HIV test and/or a seroconversion illness within the 12 months preceding HIV diagnosis.

HIV diagnosis —by exposure category and year, Victoria



HPC = High Prevalence Country (include those countries in sub-Saharan Africa, Cambodia, Myanmar, and certain countries within the Caribbean where the adult HIV prevalence is >1% and HIV is transmitted predominantly by heterosexual contact).

HIV diagnosis —2000-2004, by age group and sex, Victoria




Note: Graph excludes six transgender individuals

HIV diagnosis —2000-2004, by age group and sex, Victoria

Age Group (years)	Males		Females		Transgender	
	n	%	n	%	n	%
0-12	1	0.1	3	2.7	0	0.0
13-19	18	1.8	3	2.7	0	0.0
20-29	220	22.6	38	34.2	1	16.7
30-39	412	42.3	44	39.6	4	66.7
40-49	210	21.6	16	14.4	0	0.0
50-59	82	8.4	3	2.7	1	16.7
60+	30	3.1	4	3.6	0	0.0
Total	973	100.0	111	100.0	6	100.0

HIV diagnosis —2000-2004, by exposure category, Victoria



HPC = High Prevalence Country (include those countries in sub-Saharan Africa, Cambodia, Myanmar, and certain countries within the Caribbean where the adult HIV prevalence is >1% and HIV is transmitted predominantly by heterosexual contact).

HIV diagnosis —2000-2004, by exposure category, Victoria

Exposure Category	Males		Females		Transgender	
	n	%	n	%	n	%
Male Homosexual	748	76.9	—	—	4	66.7
Male Homosexual and Injecting Drug Use	45	4.6	—	—	2	33.3
Injecting drug use only	38	3.9	6	5.4	0	0.0
Heterosexual Contact	78	8.0	55	49.5	0	0.0
Heterosexual Contact (HPC)	45	4.6	47	42.3	0	0.0
Other/Unavailable	19	2.0	3	2.7	0	0.0
Total	973	100.0	111	100.0	6	100.0


HIV diagnosis —2000-2004, males, by heterosexual exposure category, Victoria



HPC = High Prevalence Country (include those countries in sub-Saharan Africa, Cambodia, Myanmar, and certain countries within the Caribbean where the adult HIV prevalence is >1% and HIV is transmitted predominantly by heterosexual contact).

HIV diagnosis


—2000-2004, females, by heterosexual exposure category, Victoria



HPC = High Prevalence Country (include those countries in sub-Saharan Africa, Cambodia, Myanmar, and certain countries within the Caribbean where the adult HIV prevalence is >1% and HIV is transmitted predominantly by heterosexual contact).

HIV diagnosis

—2000-2004, time between HIV diagnosis and negative test and/or seroconversion illness, by exposure category, Victoria



NPNT = No Previous Negative Test

Positive = Previously Diagnosed Positive Interstate or Overseas

HPC = High Prevalence Country (include those countries in sub-Saharan Africa, Cambodia, Myanmar, and certain countries within the Caribbean where the adult HIV prevalence is >1% and HIV is transmitted predominantly by heterosexual contact).