

## 6 Gastroenteritis

**Note for microwave oven users**

Part of the microwave cooking process includes standing time. When using a microwave, read the manufacturer's instructions carefully and observe these standing times to ensure the food is completely cooked before it is eaten.

**Household cleaning**

Bathrooms and toilets must be cleaned often to avoid the spread of germs. Pay particular attention to surfaces such as toilet seats and handles, taps and nappy change tables.

Sandpits can become contaminated with animal faeces and urine. Rake the sand frequently and remove any animal faeces. Cover the area when not in use.

**Water from untreated sources**

Untreated water that comes directly from lakes or rivers may be contaminated with faeces from people or animals. Boil water from these sources before drinking it.

**Child care centres**

Children are particularly susceptible to certain types of gastro. Nappy changing and children's lack of proper hygiene makes the transmission of these diseases in child care settings particularly high. It is important that thorough hand washing and cleaning procedures are carried out in these centres to control the spread of germs.

## 7 Gastroenteritis

**Overseas travel**

Contaminated food and drink are the most common sources of gastro in travellers, particularly those travelling to developing countries. Careful selection and preparation of food and drink offer the best protection. The main types of food and drink to avoid are uncooked food, non-bottled drinks and unpasteurised milk. Generally, cooked food that is hot, which has been thoroughly and freshly cooked, and fruit and vegetables that can be peeled or shelled, are safe.

Drinking water should be boiled or chemically treated if its purity is in doubt. Ice should be avoided.

**Note:** If you think the source of your illness may have been food, it is important that you report it to your local council health department as soon as possible. Retain (and refrigerate) any left over food which you believe may have caused you to become ill.

**Further information**

Department of Human Services  
Communicable Disease Prevention and Control Unit  
Phone: 1300 651 160  
<http://www.health.vic.gov.au/ideas>

Published by the Communicable Disease Prevention and Control Unit  
Victorian Government Department of Human Services  
Melbourne Victoria

December 2008

© Copyright State of Victoria, Department of Human Services 2004

This publication is copyright. No part may be reproduced by any process except in accordance with the provisions of the *Copyright Act 1968*.

Authorised by the State Government of Victoria, 50 Lonsdale Street,  
Melbourne.

Printed by

# Gastroenteritis



Communicable Disease Prevention  
and Control Unit  
Department of Human Services

## Gastroenteritis

### Gastroenteritis (gastro) – what is it and how did I get it?

Gastro is an illness which may cause some or all of the following symptoms:

- diarrhoea
- stomach cramps
- vomiting
- nausea
- fever
- headache

Germs which cause gastro get into your body through your mouth and it usually takes several days for symptoms to appear. This time is known as the incubation period. Sometimes symptoms may occur within hours of the germs entering your body but it is usually longer.

When people get gastro they often assume that the last meal they ate gave them food poisoning. This is usually not the case.

### What causes gastro and how is it spread?

The most common germs that cause gastroenteritis are bacteria, viruses and certain parasites. They may be found in soil, wild and pet animals including birds, and humans.

Gastro occurs when these germs are taken in by mouth and this may happen in any of the following ways:

- from person to person – this may occur directly by close personal contact or contact with the faeces of an infected person, or indirectly by touching

contaminated surfaces such as taps, toilet flush handles, children's toys and nappies

- eating contaminated food
- drinking contaminated water
- airborne through vomiting, coughing and sneezing (mainly viruses).
- handling pets and other animals

People and animals can carry the germs which cause gastro in their faeces without having any symptoms. These people and animals can still pass the disease on to others.

### How can food and water become contaminated?

Food can become contaminated by people who have gastro, if they do not wash their hands properly after going to the toilet and before handling food.

Bacteria which can cause gastro are often present in raw foods such as meats, poultry and eggs. These raw foods must always be handled, prepared and stored so as not to contaminate other foods. Proper cooking will kill these bacteria.

If insects, rodents or other animals are not stopped from entering areas where food is prepared, they may contaminate food, equipment, benches and utensils with gastroenteritis germs.

Creeks, rivers, lakes and dams may be polluted with faeces from humans or animals. You should not drink water from these sources, unless it has been treated or boiled.

### I think I may have gastroenteritis – what should I do?

If you have symptoms of gastroenteritis, report them to your doctor. This will ensure that you receive proper treatment and advice and that steps are taken to reduce the spread of the disease.

### Can I still work?

As gastro is very infectious, it is advisable that people with symptoms should not work or attend childcare centres, kindergartens or schools until 48 hours after symptoms have stopped.

It is particularly important that food handlers, child care workers and health care workers do not work for at least 48 hours after symptoms have stopped.

### How can I avoid getting gastro?

In your household, the risk of spreading gastro can be reduced. It is very important that people with gastro do not prepare or handle food that is to be eaten by other people and that no one shares their towel, face washer, toothbrush or eating utensils.

By following the guidelines below, everyone can do something to avoid getting gastro.

### Careful hand washing

Everyone should wash their hands thoroughly with soap and warm running water for at least 15 seconds:

- before preparing food
- between handling raw and ready-to-eat foods

- before eating
- after going to the toilet or changing nappies
- after smoking
- after using a tissue or handkerchief
- after working in the garden
- after playing with pets

Food handlers should use disposable paper towels or an air dryer to dry their hands. Cloth towels are not recommended as they get dirty quickly and can spread germs from one person to another.

### Safe food storage and handling

- Do not handle raw and cooked foods with the same implements (tongs, knives, cutting boards), unless they have been thoroughly washed between uses
- Keep all kitchen surfaces and equipment clean
- Defrost food by placing it on the lower shelves of a refrigerator or use a microwave oven
- Thoroughly cook all raw foods
- Place cooked food in a refrigerator within an hour of cooking it
- Refrigerate raw foods below cooked or ready-to-eat foods to prevent cross-contamination
- Store food below 5°C or above 60°C to prevent growth of germs
- Thoroughly wash raw vegetables before eating
- Reheat food until the internal temperature of the food reaches at least 75°C
- Protect food from insects, rodents and other animals