

# Recommended Immunisation Schedule



AGE	IMMUNISATION
The following immunisations are recommended at the stated age.	
Birth	Hepatitis B Vaccine
2 months	Diphtheria/Tetanus/Pertussis Vaccine Oral Polio Vaccine Hib (Haemophilus influenzae type b)/ Hepatitis B Vaccine <i>Childhood Pneumococcal Vaccine*</i>
4 months	Diphtheria/Tetanus/Pertussis Vaccine Oral Polio Vaccine Hib (Haemophilus influenzae type b)/ Hepatitis B Vaccine <i>Childhood Pneumococcal Vaccine*</i>
6 months	Diphtheria/Tetanus/Pertussis Vaccine Oral Polio Vaccine <i>Childhood Pneumococcal Vaccine*</i>
12 months	Measles/Mumps/Rubella Vaccine Hib (Haemophilus influenzae type b)/ Hepatitis B Vaccine Meningococcal C Vaccine
18 months	Diphtheria/Tetanus/Pertussis Vaccine
4 years	Diphtheria/Tetanus/Pertussis Vaccine Oral Polio Vaccine Measles/Mumps/Rubella Vaccine
Year 7 (secondary school)	Hepatitis B Vaccine (1st dose) Hepatitis B Vaccine (2nd dose) 4 – 6 months apart
Year 10 (secondary school)	Adult Diphtheria/Tetanus Vaccine
Non-immune women shortly after delivery	Measles/Mumps/Rubella Vaccine
50 years	Adult Diphtheria/Tetanus Vaccine
Over 50 years (Aboriginal & Torres Strait Islander people)	Influenza Vaccine (annual) Pneumococcal Vaccine (5 yearly)
Over 65 years	Influenza Vaccine (annual) Pneumococcal Vaccine (5 yearly)

*\*Funded for Aboriginal and Torres Strait Islanders up to 2 years of age and children with certain medical risk factors as recommended by a Paediatrician.*

We gratefully acknowledge the contribution of the Department of Health and Ageing for this information.

