

Croup

What is croup?

Croup (medically called *laryngotracheitis*) is a viral infection of the throat, which affects the voice box (*larynx*) and windpipe (*trachea*).

Croup is common in babies and young children during winter. Most cases are mild. These include cases where a child has a 'barking' cough, but no trouble breathing. Children are best cared for at home.

Sometimes croup can become more serious with the child having difficulty breathing.

What causes croup?

Children have small, soft windpipes. A viral infection causes the lining of the windpipe and voice box to swell. This, along with a build-up of sticky fluid (*mucous*), narrows the passageway through which air travels into the lungs. This can make it difficult for your child to breathe.

As children get older, the size and strength of the windpipe improves, which is why croup is less common in older children.

What are the symptoms of croup?

Croup usually starts as a common cold (with a runny nose, red eyes, sore throat and a fever). Then the following symptoms develop:

- a hoarse voice
- a harsh, 'barking' cough (like a seal's bark)
- noisy breathing (a high pitched noise when breathing in, known as *stridor*)
- fast or laboured (hard) breathing.

These symptoms often start at night when the air is cold and dry. They can get worse when your child is upset.

Treatment

- In the emergency department, your child may receive a medicine called *prednisolone* or *dexamethasone*. This helps reduce the swelling in the airways, making it easier for your child to breathe. It will work in four to six hours.
- If the croup is severe, another medicine, called *adrenaline*, may be given through a nebuliser (which lets you breathe in the medication) or an injection. This relieves the spasm and swelling until the *prednisolone* starts to work.
- **Antibiotics will not help**, as croup is caused by a virus.
- There is no proof that anti-allergy medication (*antihistamines*) or cold remedies (*decongestants*) are of benefit in croup.

Home care

- Stay calm. Try to keep your child calm also. Croup can be very frightening for a child, but being scared makes it worse. Try watching television or reading a book with your child to help them relax.
- Offer your child plenty to drink. Offer small amounts ('sips') and often, especially if your child has a fever. Water is best.
- Sit your child up in bed with pillows to make breathing easier.
- Offer paracetamol (such as *Panadol* or *Dymadon*) if your child has a fever and is miserable. Carefully check the label for the correct dose and make sure you are not giving your child any other products containing paracetamol (such as some cough medicines and cold and flu preparations).
- Steam therapy does not help. It is also regarded as dangerous because of accidental burns, so is not advised.

What to expect

- Croup may last one or two days and only occur at night. Sometimes it can last up to a week, and may be present during the day as well (it is usually worse at night). The first two days are usually the worst.
- Croup is often a mild illness but can get worse quickly. Do not hesitate to seek medical help if this is the case. A child with severe croup will need prompt treatment in hospital.
- Some children can get croup more than once. It is not possible to prevent croup.



Croup

Notes:

Seeking help



In a **medical emergency** go to the nearest hospital emergency department or call an ambulance (dial 000). Do this if your child:

- has trouble breathing
- becomes floppy or hard to wake
- goes pale or blue in colour
- sucks in their breastbone or ribcage when breathing
- looks very sick and is dribbling
- is very distressed
- has *stridor* (noisy breathing) **at rest**.



For other medical problems see your local doctor or health care professional.



For health advice from a Registered Nurse you can call **NURSE-ON-CALL 24 hours a day** on **1300 60 60 24** for the cost of a local call from anywhere in Victoria.*

NURSE-ON-CALL provides access to **interpreting services** for callers not confident with English. Call **1300 60 60 24**.

*Calls from mobile calls may be charged at a higher rate



Want to know more?

- Ask your local doctor or health care professional.
- Visit the **Royal Children's Hospital** website www.rch.org.au/kidsinfo
- Visit the **Better Health Channel** www.betterhealth.vic.gov.au

If you would like to receive this publication in an **accessible format**, please phone 9096 8064 or email edfactsheets@dhs.vic.gov.au

June 2008. Also available online at www.health.vic.gov.au/edfactsheets

Disclaimer: This health information is for general education purposes only. Please consult with your doctor or other health professional to make sure this information is right for you.