Immunisation schedule Victoria April 2025

This schedule is intended as a guide only and complete consultation with the Australian Immunisation Handbook is recommended.

National Immunisation Schedule - Infants and children

pertussis-poliomyelitis-hepatitis B-Haemophilus Influenzae type b 6 months to <5 years (annually) 12 months Measles-mumps-rubella Priorix or M-M-R II (IM/SC) Meningococcal ACWY Pneumococcal B Measles-mumps-rubella Aboriginal* Measles-mumps-rubella Diphtheria-tetanus-pertussis pertussis poliomyelitis Type b As supplied Administer annually, In children aged 6 months to less than 9 years of age in the first year of administration, give 2 doses a minimum of 1 month apart Premature babies < 32 weeks gestation or <2000g birthweight are recommended to receive an additional dose of H-B-Vax-II padiatric or Engerix-B paddiatric – see AlH® Bexsero (IM) Aboriginal* Meningococcal B Bexsero (IM) Priorix-tetra (IM/SC) Tubella-varicella Diphtheria-tetanus-pertussis Infanrix or Tripacel (IM) Haemophilus Influenzae type b At-HIB (IM/SC) Quadracel (IM) Children with at risk conditions for IPD* are recommended to receive an additional dose of Pneumovax 23 – see AlH® Administer annually. If less than 9 years of age in the first year of administration, give 2 doses a minimum of 1 month apart Aboriginal* ≥ 5 years (annually) Influenza† As supplied Administer annually. If less than 9 years of age in the first year of administration, give 2 doses a minimum of 1 month apart Administer annually. If less than 9 years of age in the first year of administration, give 2	National inimumsation schedule -		illiants and ciliuren		
2 months (from 6 weeks) and pertussis, poliomyelitishepatitis B-Hoemophilus Influenzae type b Pneumococcal Prevenar 13 (IM) Pneumococcal B Prevenar 13 (IM) Prevenatible B-Hoemophilus Influenzae type b Pneumococcal B Prevenar 13 (IM) Poartis (IM) Poartis (IM) Poartis (IM) Poartis (IM) Poartis (IM) Prevenar 13 (IM) Poartis (IM) Prevenar 13 (IM) Poartis (IM) Poarti	Age	Disease	Vaccine	Notes	
and A pertussis, poliomyelitis- hepatitis B-Haemophillus Influenzae type b Pneumococcal Prevenar 13 (IM) Rotarivus Rotarix (Oral) Rotariv (Oral) Rotarivis (Oral) Resserce (IM) (IM) or Vacelis (IM) or Vacelis (IM) or Vacelis (IM) Influenzae type b Rotarivis (Oral) Rotarivis (Oral) Resserce (IM) (Im) or Capture 26 months with at risk conditions for IPD* are recommended to receive an additional dose of Prevenar 13 – see AlH* (IM) or IPD* are recommended to receive an additional dose of Prevenar 13 – see AlH* (IM) or IPD* are recommended to receive an additional dose of Prevenar 13 – see AlH* (IM) or IPD* are recommended to receive an additional dose of Prevenar 13 – see AlH* (IM) or IPD* are recommended to receive an additional dose of Prevenar 13 – see AlH* (IM) or IPD* are recommended to receive an additional dose of Prevenar 13 (IM) Realize - Turning	Birth	Hepatitis B	or Engerix-B		
Pneumococcal Prevenar 13 (IM) dose 2 limited to 10-24 weeks of age Bexsero: Prophylactic paracetamol recommended pre and post vaccination – see AIH*. Catch up available for Aboriginal* As supplied Administer annually. In children ≥ years Infanrix hexo (IM) or Vaxelis (IM) Premature basics ≥ 32 weeks gestation or excelled a minimum of 1 month apart Premature basics ≥ 32 weeks gestation or excelled a minimum of 1 month apart Priorix-ctera (IM) ≤ 5 years Infanrix hexo (IM) Premature basics ≥ 32 weeks gestation or excelled a minimum of 1 month apart Premature basics ≥ 32 weeks gestation or excelled a minimum of 1 month apart Premature basics ≥ 32 weeks gestation or excelled a minimum of 1 month apart Premature basics ≥ 32 weeks gestation or excelled a minimum of 1 month apart Premature basics ≥ 32 weeks gestation or excelled a minimum of 1 month apart Premature basics ≥ 32 weeks gestation or excelled a minimum of 1 month apart Premature basics ≥ 32 weeks gestation or excelled a minimum of 1 month apart Premature basics ≥ 32 weeks gestation or excelled a minimum of 1 month apart Premature basics ≥ 32 weeks gestation or excelled a minimum of 1 month apart Premature basics ≥ 32 weeks gestation or excelled a minimum of 1 month apart Premature basics ≥ 32 weeks gestation or excelled a minimum of 1 month apart Premature basics ≥ 32 weeks gestation or excelled a minimum of 1 month apart Premature basics ≥ 32 weeks gestation or excelled a minimum of 1 month apart Premature basics ≥ 32 weeks gestation or excelled a minimum of 1 month apart Premature basics ≥ 32 weeks gestation or excelled a minimum of 1 month apart Premature basics ≥ 32 weeks gestation or excelled a minimum of 1 month apart Premature basics ≥ 32 weeks gestation or excelled a minimum of 1 month apart Premature basics ≥ 32 weeks gestation or excelled a minimum of 1 month apart Premature basics ≥ 32 weeks gestation or excelled a minimum of 1 month apart Premature basics ≥ 32 weeks gestation or exce	and	pertussis, poliomyelitis- hepatitis B- <i>Haemophilus</i>	` '	Rotarix: Dose 1 limited to 6-14 weeks of age.	
Aboriginal [‡] As supplied Administer annually. If less than 9 years of age in the first year of administration, give 2		Pneumococcal	Prevenar 13 (IM)	dose 2 limited to 10-24 weeks of age	
Aborginal* Meningococcal B Bessero (IM) children <2 years 6 months Diphtheria-tetanus-pertussis-poliomyelitis hepatitis B-Haemophilus Influenzae type b Infanrix hexa (IM) or Vaxelis (IM) Children ≥6 months with at risk conditions for IPD* are recommended to receive an additional dose of Prevenar 13 – see AlH* 6 months to <5 years (annually)		Rotavirus	Rotarix (Oral)	recommended pre and post vaccination – see	
Pertussis-pollomyelitis-hepatitis B-Haemophilus Influenzae type b Promise type type b Promise type type b Promise type type type type type type type typ	Aboriginal [‡]	Meningococcal B	Bexsero (IM)		
Cannually Cann	6 months	pertussis-poliomyelitis- hepatitis B- <i>Haemophilus</i>	` '		
Meningococcal ACWY Nimenrix (IM) Pneumococcal ACWY Prevenar 13 (IM) Pneumococcal Prevenar 13 (IM) Pneumococcal Prevenar 13 (IM) Pneumococcal Prevenar 13 (IM) Pneumococcal Prevenar 13 (IM) Pneumococcal Prevenar 13 (IM) Pneumococcal Prevenar 13 (IM) Pneumococcal Prevenar 13 (IM) Pneumococcal Prevenar 13 (IM) Pneumococcal Prevenar 13 (IM) Prevenar 13 (IM) Pneumococcal Pne		Influenza†	As supplied	months to less than 9 years of age in the first year of administration, give 2 doses a	
Meningococcal ACWY Nimenrix (IM) paediatric or Engerix-B paediatric – see AIH#		Measles-mumps-rubella		<2000g birthweight are recommended to receive an additional dose of H-B-Vax-II paediatric or Engerix-B paediatric – see AIH# Bexsero: Prophylactic paracetamol recommended pre and post vaccination – see AIH#. Catch up available for Aboriginal‡	
Aboriginal [‡] Meningococcal B Bexsero (IM) Mesles-mumps-rubella-varicella Diphtheria-tetanus-pertussis (IM) Haemophilus Influenzae type b Act-HIB (IM/SC) Tiphtheria-tetanus-pertussis-poliomyelitis Diphtheria-tetanus-pertussis-poliomyelitis As supplied As supplied Administer annually. If less than 9 years of age in the first year of administration, give 2 doses a minimum of 1 month apart Aboriginal [‡] ≥ 5 years (annually) Measles-mumps-rubella Bexsero (IM) Priorix-tetra (IM/SC) Infanrix or Tripacel (IM) Act-HIB (IM/SC) Children with at risk conditions for IPD* are recommended to receive an additional dose of Pneumovax 23 – see AIH [#] As supplied Administer annually. If less than 9 years of age in the first year of administration, give 2 doses a minimum of 1 month apart Aboriginal [‡] ≥ 5 years (annually) Influenza [†] As supplied Administer annually. If less than 9 years of age in the first year of administration, give 2		Meningococcal ACWY	Nimenrix (IM)		
thildren <2 years Measles-mumps-rubella-varicella Diphtheria-tetanus-pertussis Infanrix or Tripacel (IM)		Pneumococcal	Prevenar 13 (IM)		
rubella-varicella Diphtheria-tetanus- pertussis		Meningococcal B	Bexsero (IM)		
pertussis (IM) Haemophilus Influenzae type b Act-HIB (IM/SC) 4 years Diphtheria-tetanus-pertussis-poliomyelitis Quadracel (IM) Style b Children with at risk conditions for IPD* are recommended to receive an additional dose of Pneumovax 23 − see AIH# Style 5 years with at risk condition (annually) Influenza† As supplied Administer annually. If less than 9 years of age in the first year of administration, give 2 doses a minimum of 1 month apart Aboriginal ≥ 5 years (annually) Influenza† As supplied Administer annually. If less than 9 years of age in the first year of administration, give 2	18 months	· · · · · · · · · · · · · · · · · · ·	Priorix-tetra (IM/SC)		
type b 4 years Diphtheria-tetanus-pertussis-poliomyelitis Syears with at risk conditions for IPD* are recommended to receive an additional dose of Pneumovax 23 − see AIH# Syears with at risk condition (annually) As supplied As supplied Administer annually. If less than 9 years of age in the first year of administration, give 2 doses a minimum of 1 month apart Aboriginal* Syears (annually) As supplied Administer annually. If less than 9 years of age in the first year of administration, give 2 doses a minimum of 1 month apart		•			
pertussis-poliomyelitis Quadracel (IM) recommended to receive an additional dose of Pneumovax 23 − see AIH [#] ≥ 5 years with at risk condition (annually) Influenza [†] As supplied Administer annually. If less than 9 years of age in the first year of administration, give 2 doses a minimum of 1 month apart Aboriginal [‡] ≥ 5 years (annually) As supplied Administer annually. If less than 9 years of age in the first year of administration, give 2		· · · · · · · · · · · · · · · · · · ·	Act-HIB (IM/SC)		
condition (annually) age in the first year of administration, give 2 doses a minimum of 1 month apart Aboriginal [‡] ≥ 5 years (annually) Influenza [†] As supplied age in the first year of administration, give 2	4 years	-		recommended to receive an additional dose	
(annually) age in the first year of administration, give 2	-	Influenza†	As supplied	age in the first year of administration, give 2	
		Influenza†	As supplied		

National Immunisation Schedule - Adolescents, adults and at-risk groups

Age	Disease	Vaccine	Notes	
Aboriginal [‡] (annually)	Influenza†	As supplied	Administer annually	
People with at risk conditions (annually)	Influenza†	As supplied	Administer annually	
Year 7 (or age equivalent)	Human papillomavirus (HPV)	Gardasil 9 (IM)	HPV: Single dose recommended. Not required if previously received. If dose was missed and not previously received, a catch-up is available up to and including 25 years of age	
	Diphtheria-tetanus- pertussis	Adacel or Boostrix (IM)		
Year 10 (or age equivalent)	Meningococcal ACWY	MenQuadfi (IM)		
People with at risk conditions for IPD*	Pneumococcal	Prevenar 13 and Pneumovax 23 (IM)	See AIH# for conditions recommended to receive Prevenar 13 and Pneumovax 23	
All people with asplenia, hyposplenia, complement deficiency and treatment	Meningococcal ACWY	Nimenrix (IM)	See AIH# for required doses and timing Additional groups are	
with eculizumab	Meningococcal B	Bexsero (IM)	recommended to receive these vaccines but these are not funded	
>5 years with asplenia or hyposplenia	Haemophilus Influenzae type b (HiB)	Act-HIB (IM)	If incompletely vaccinated or not vaccinated in childhood	
Eligible people 18 years and over considered at increased risk of herpes zoster due to an underlying condition and/or immunomodulatory/immunosuppressive treatments	Shingles (Herpes Zoster)	Shingrix (IM)	See AIH# for required doses and timing See Further Information section for shingles vaccination eligibility	
All people under 20 years of age	Age-appropriate vaccine catch-up	As supplied	Refugees and humanitarian entrants of any age are eligible for free catch-up vaccines	
Pregnant	Influenza†	As supplied	Influenza†: Any trimester	
	Pertussis (whooping cough)	Adacel or Boostrix (IM)	Pertussis: Each pregnancy between 20-32 weeks gestation	
	Respiratory Syncytial Virus (RSV)	Abrysvo (IM)	RSV: 28-36 weeks gestation	
Aboriginal [‡] ≥50 years	Pneumococcal	Prevenar 13 then Pneumovax 23 (IM)	See AIH [#] for required doses and timing	
	Shingles (Herpes Zoster)	Shingrix (IM)		
≥ 65 years	Influenza†	As supplied	Administer annually	
≥ 65 years	Shingles (Herpes Zoster)	Shingrix (IM)	See AIH# for required doses and timing	
From 70 years	Pneumococcal	Prevenar 13 (IM)		

- The <u>National Immunisation Program (NIP)</u> provides the above routine vaccinations free to infants, children, adolescents and adults who have, or are eligible for a Medicare benefits.
- All people less than 20 years of age are eligible for NIP vaccines missed in childhood, except for HPV which is available free up to and including age 25.
- The number and range of vaccines and doses that are eligible for NIP funded catch-up is different for people aged less than 10 years and those aged 10–19.
- All refugees and humanitarian entrants of any age are eligible for the following vaccines if they were missed: diphtheria-tetanus-pertussis, chickenpox, poliomyelitis, measles-mumps-rubella and hepatitis B as well as HPV (up to and including age 25). Refer to the Immunisation Handbook for timing of doses.

Victorian Government funded immunisation program

The below vaccines and immunisation products are funded by the Victorian Government. See Immunisation schedule Victoria and vaccine eligibility criteria online for further information.

Disease	Vaccine	Age/at risk conditions
Tuberculosis (BCG)	As supplied (ID)	 BCG for infants and children under 5 years of age travelling to TB endemic countries. See AIH# for BCG dosing and vaccination procedures.
Measles-mumps- rubella	As supplied (IM/SC)	 Infants from 6 months and before 11 months of age prior to overseas travel. This dose is in addition to the scheduled MMR vaccine doses recommended at ages 12 (MMR) and 18 months (MMRV) Women planning pregnancy or after delivery
		People born during or since 1966
Age-appropriate vaccine catch-up	As supplied	Vulnerable citizens – a person who has experienced socioeconomic disadvantage that compromised their equitable access to the vaccine during their period of eligibility. Persons are vaccinated based on an individual assessment by an immunisation provider. See Immunisation schedule Victoria and vaccine eligibility criteria online for more information
Hepatitis B	As supplied (paediatric or adult) (IM)	Eligible people at risk - see Immunisation schedule Victoria and vaccine eligibility criteria online for more information
Rabies	As supplied	Australian volunteer wildlife handlers working in a volunteer service in Victoria for the initial course of pre-exposure rabies vaccine
		Post exposure treatment for rabies or Australian Bat Lyssavirus
		 See Immunisation schedule Victoria and vaccine eligibility criteria online for more information
Respiratory Syncytial Virus (RSV)	Immunisation product - Monoclonal antibody Beyfortus™ (nirsevimab) (IM)	 Infants born between 1 April - 30 September 2025 to mothers who did not receive maternal RSV vaccine during pregnancy, or where maternal RSV vaccine was administered less than 2 weeks before birth or with risk conditions for severe RSV disease
		 1st RSV season - Infants born from 1 October 2024 - 31 March 2025 to mothers who did not receive maternal RSV vaccine during pregnancy, or where maternal RSV vaccine was administered less than 2 weeks before birth or with risk conditions for severe RSV disease
		 2nd RSV season -Infants born on or after 1 October 2023 vulnerable to severe RSV disease including Aboriginal[‡] and young children with increased risk of severe RSV disease
		 For more information see RSV immunisation and the #AIH RSV section for risk conditions for RSV disease

*IPD: Invasive pneumococcal disease. #AIH: Online Australian Immunisation Handbook. †Influenza: For information on age-appropriate vaccines or specified medical risk conditions refer to the AIH# or the annual ATAGI advice on seasonal Influenza vaccines. ‡Aboriginal: The term Aboriginal is inclusive of Aboriginal and Torres Strait Island people.

Further information

The Australian Immunisation Handbook: https://immunisationhandbook.health.gov.au

Immunisation schedule Victoria and vaccine eligibility criteria: https://www.health.vic.gov.au/immunisation/immunisation-rehadule victoria and vaccine eligibility criteria

schedule-victoria-and-vaccine-eligibility-criteria

Shingles vaccination: https://www.health.gov.au/resources/publications/national-immunisation-program-shingles-

vaccination-program-advice-for-health-professionals-september-2024?language=en

RSV immunisation: https://www.health.vic.gov.au/immunisation/respiratory-syncytial-virus-immunisation

Ordering vaccines: https://www.health.vic.gov.au/immunisation/ordering-vaccines

Cold chain management: https://www.health.vic.gov.au/immunisation/cold-chain-management

Australian Immunisation Register: 1800 653 809

SAEFVIC - Victoria's vaccine safety service: 1300 882 924 #1

To receive this document in another format, email Immunisation Program immunisation@health.vic.gov.au.

Authorised and published by the Victorian Government, 1 Treasury Place, Melbourne. © State of Victoria, Australia, Department of Health, April 2025.

ISBN 978-1-76131-757-6 (pdf/online/MS word)

Available at Immunisation schedule Victoria and vaccine eligibility criteria https://www.health.vic.gov.au/immunisation/immunisation-schedule-victoria-and-vaccine-eligibility-criteria >