



# Victorian Respiratory Surveillance Report

23 January 2026

## Contents

<b>About this report</b>	<b>1</b>
<b>Summary</b>	<b>2</b>
<b>Laboratory surveillance</b>	<b>3</b>
Case notifications	3
Testing	9
<b>Community surveillance</b>	<b>12</b>
FluTracking	12
<b>How to use this report</b>	<b>13</b>
Data sources	13
Definitions	13

## About this report

The Victorian Respiratory Surveillance Report summarises the latest surveillance information on COVID-19, influenza and respiratory syncytial virus (RSV) in Victoria. This report provides an overview of the current levels and trends of acute respiratory illness from these conditions in the community. Please see the section on How to use this report for more information.

Measurements of the greatest value for summarising respiratory illness in Victoria are presented in this report. Other systems are also reviewed by the Department of Health to understand patterns of respiratory disease in Victoria which inform the overall summaries presented.

Data are presented as at 21 January 2026 for the week ending 17 January 2026.

Each report reflects the data available at the time the report was prepared. Information may differ from previous reports as new data are received or updated.



Department of Health

# Summary

In Victoria, activity of COVID-19 and RSV are at inter-seasonal levels, and influenza has continued to decrease.


## COVID-19 activity is low

### CASE TRENDS (LAST 2 WEEKS)

Notifications have decreased



### NOTIFICATIONS LAST 12 MONTHS



### TEST POSITIVITY (LAST 2 WEEKS)

The percentage of tests that were positive have decreased



COVID-19 activity has remained low following the winter wave.

## Influenza activity is low-moderate

### CASE TRENDS (LAST 2 WEEKS)

Notifications have decreased



### NOTIFICATIONS LAST 12 MONTHS



### TEST POSITIVITY (LAST 2 WEEKS)

The percentage of tests that were positive have continued to decrease



Influenza activity has decreased in Victoria following a prolonged season.

## RSV activity is low

### CASE TRENDS (LAST 2 WEEKS)

Notifications have decreased



### NOTIFICATIONS LAST 12 MONTHS



### TEST POSITIVITY (LAST 2 WEEKS)

The percentage of tests that were positive have slightly decreased



RSV activity has returned to typical inter-seasonal levels for this time of year.

# Laboratory surveillance

## Case notifications

Laboratory-confirmed cases of COVID-19, influenza and RSV are notified to the Victorian Department of Health. Notified infections that are diagnosed through laboratory testing are only a subset of the total number of infections in the community. Trends in notifications may be impacted by changes in testing.


### Notifications (last two weeks)

In the past week, COVID-19 notifications have decreased (-37%), influenza notifications have decreased (-45%) and RSV notifications have decreased (-38%).



## COVID-19

**Figure 1:** COVID-19 notified cases by week, Victoria, 19 January 2025 to 17 January 2026



There were **389** notified COVID-19 cases two weeks ago


4 Jan 2026 to 10 Jan 2026

There were **246** notified COVID-19 cases last week

11 Jan 2026 to 17 Jan 2026

**37%** decrease

**Figure 2:** COVID-19 trends in notified cases (7-day rolling average), Victoria, 1 January 2023 to 17 January 2026




Notified COVID-19 cases are lower than the same time in the past three years

## Influenza

Influenza notifications presented in this report are reported from a subset of laboratories in Victoria, generally comprising around 85% of total influenza notifications.

**Figure 3:** Influenza notified cases by week, Victoria, 19 January 2025 to 17 January 2026



There were **999** cases of influenza (overall) notified two weeks ago

4 Jan 2026 to 10 Jan 2026

There were **550** cases of influenza (overall) notified last week

11 Jan 2026 to 17 Jan 2026



**45%** decrease

There were **976** cases of influenza A notified two weeks ago

4 Jan 2026 to 10 Jan 2026

There were **531** cases of influenza A notified last week

11 Jan 2026 to 17 Jan 2026



**46%** decrease

There were **23** cases of influenza B notified two weeks ago

4 Jan 2026 to 10 Jan 2026


There were **19** cases of influenza B notified last week

11 Jan 2026 to 17 Jan 2026




**17%** decrease

**Figure 4:** Influenza A trends in notified cases (7-day rolling average), Victoria, 1 January 2023 to 17 January 2026



Notified influenza A cases are similar to the same time in the past three years

**Figure 5:** Influenza B trends in notified cases (7-day rolling average), Victoria, 1 January 2023 to 17 January 2026




Notified influenza B cases are at inter-seasonal levels

## Respiratory Syncytial virus (RSV)

RSV notifications presented in this report are reported from a subset of laboratories in Victoria, generally comprising around 85% of total RSV notifications.

**Figure 6:** RSV notified cases by week, Victoria, 19 January 2025 to 17 January 2026



There were **187** notified RSV cases two weeks ago


4 Jan 2026 to 10 Jan 2026

There were **116** notified RSV cases last week

11 Jan 2026 to 17 Jan 2026

**38%** decrease

**Figure 7:** RSV trends in notified cases (7-day rolling average), Victoria, 1 January 2023 to 17 January 2026




Notified RSV cases are at inter-seasonal levels

## Young children and older adults


Young children and older adults are at greater risk of severe disease from RSV.

**Figure 8:** RSV trends in notified cases by week in young children (under 5 years), Victoria, 19 January 2025 to 17 January 2026



**i** Children under 2 years comprise the majority of notified RSV cases in young children

**Figure 9:** RSV trends in notified cases by week in older adults (65+ years), Victoria, 19 January 2025 to 17 January 2026



**i** Notified RSV cases in adults aged over 65 years are at inter-seasonal levels


# Testing

Laboratory testing for respiratory illnesses changes over time. Tracking the percentage of tests with the notifiable condition detected (i.e. test positivity) is a useful measure to understand trends in disease surveillance over time.


Test results presented in this report are from selected laboratories. These include private and hospital laboratories and represent tests completed across Victoria.

## Summary

In the past week, the percentage of COVID-19 tests that were positive have decreased (2.9% to 2.2%), the percentage of influenza tests that were positive have continued to decrease (4.8% to 3.4%) (influenza A: 9.3% to 6.5%; influenza B: 0.2% to 0.2%), and the percentage of RSV tests that were positive have slightly decreased (1.8% to 1.4%).




**Figure 10:** Weekly test positivity for COVID-19, influenza and RSV, Victoria, 19 January 2025 to 17 January 2026



- ⓘ COVID-19 test positivity has decreased slightly in recent weeks
- ⓘ Influenza A test positivity has decreased in recent weeks
- ⓘ Influenza B test positivity is at inter-seasonal levels
- ⓘ RSV test positivity is at inter-seasonal levels

**Figure 11:** Weekly test positivity by age groups for COVID-19, influenza and RSV, Victoria, 19 January 2025 to 17 January 2026



# Community surveillance

Respiratory illnesses are not limited to the notifiable conditions presented above. Understanding the overall burden of respiratory illness in the community is useful to understand broader trends in illness over time.


## FluTracking

FluTracking is an online surveillance system across Australia, New Zealand, Hong Kong and Fiji. Volunteer participants complete a simple weekly online survey which collects self-reported information on respiratory symptoms. Data used in this report are received up to 12:00PM Tuesday. More information about FluTracking and ways to be involved are available here: <https://info.flutracking.net/>

### Summary

In the past week, the percentage of Victorian FluTracking participants reporting respiratory illness (fever and cough) were stable from the previous week.

**Figure 12:** Proportion of FluTracking participants reporting respiratory illness by week, Victoria, 1 January 2023 to 18 January 2026. Respiratory illness is defined as fever & cough.



Reported respiratory illness from FluTracking is at lower levels compared to the past three years

# How to use this report

## Data sources

All notified cases in Victoria are recorded in the Victorian Public Health Events Surveillance System (PHESS). Under the *Public Health and Wellbeing Act 2008* and *Public Health and Wellbeing Regulations 2019*, the Victorian Department of Health is authorised to collect information from doctors and laboratories about diagnoses of certain health-related conditions in Victoria.

Modelling is performed by the Australia-Aotearoa Consortium for Epidemic Forecasting & Analytics (ACEFA) throughout the winter respiratory season using data provided by the Department of Health. <https://github.com/acefa-hubs>

Pathology services are required to notify the Department of Health of all tests performed during a weekly period for schedule 4A conditions including influenza, RSV and COVID-19.

The FluTracking surveillance system collects data from volunteer participants in a weekly survey. FluTracking reports are available here: <https://info.flutracking.net/reports/australia-reports/>

The Victorian Department of Health continually reviews surveillance methods to monitor respiratory disease in Victoria. Measurements included in this report may be updated or removed accordingly.

## Definitions

**Notified cases:** Laboratory-confirmed cases of COVID-19, influenza, and RSV are reported according to the CDNA case definitions <https://www.health.gov.au/resources/collections/cdna-surveillance-case-definitions/> Where multiple positive test results are received for the same person within 30 days of the initial test result they are counted as a single case.

Rapid antigen test results are not collected by the Victorian Department of Health, however, remain an important tool for individuals to access treatment and protect their community.

Case notifications are included in the Victorian dataset if the postcode of residence of the case is in Victoria. This does not necessarily reflect where the infection was acquired.

**Modelled current trends:** Modelled current trends indicate whether cases are likely to be increasing, decreasing or stable based on modelled trends in expected cases for each condition. Details of the statistical modelling methods can be found in the paper at <https://doi.org/10.1093/aje/kwaf119>

**Test positivity:** Test positivity is the percentage of total tests where the notifiable condition was detected. This is reported by the date of specimen collection.

**Dates:** Case notifications are based on the date the notification was first received by the Victorian Department of Health. Test positivity is based on the date of specimen collection.

**Weeks:** For the purposes of this report, data are aggregated by week, with the week starting Sunday and ending Saturday unless otherwise specified.

**Activity levels:** The Victorian Department of Health monitors activity of acute respiratory illnesses assessed against trends from recent seasons and determined by expert review of multiple surveillance inputs informed by statistical analysis of all data sources available at the time the report is prepared.